

Heavy metal assessment and water quality values in urban stream and rain water

¹*K. Sekabira; ²H. Oryem Origa; ³T. A. Basamba; ²G. Mutumba; ²E. Kakudidi

¹Department of Environment, Faculty of Science and Engineering, Kampala International University, Kampala, Uganda

²Department of Botany, Faculty of Science, Makerere University, Kampala, Uganda

³Department of Soil Science, Faculty of Agriculture, Makerere University, Kampala, Uganda

Received 2 April 2010; revised 14 June 2009; accepted 21 July 2010; available online 1 September 2010

ABSTRACT: Water quality monitoring in developing countries is inadequate, especially in stream water affected by urban effluents and runoff. The purpose of this study was to investigate heavy metal contaminants in the Nakivubo Stream water in Kampala, Uganda. Water samples Nakivubo Channelized Stream, tributaries and industrial effluents that drain into the stream were collected and analysed for the total elemental concentration using flame atomic absorption spectrophotometer. The results showed that: 1) the wastewater was highly enriched with lead and manganese above the maximum permissible limit; 2) the levels of dissolved oxygen were below the maximum permissible limit, while the biological oxygen demand was above the maximum permissible limit. All industrial effluents/wastewater were classified as strong (> 220 mg/L). Factor analysis results reveal two sources of pollutants; 1) mixed origin or chemical phenomena of industrial and vehicular emissions and 2) multiple origin of lead (vehicular, commercial establishment and industrial). In conclusion, Nakivubo Channelized Stream water is not enriched with heavy metals. These heavy metals (lead, cadmium and zinc) were rapidly removed by co-precipitation with manganese and iron hydroxides and total dissolved solids into stream sediments. This phenomena is controlled by pH in water.

Keywords: Co-precipitation; Factor analysis; Industrial emission; Nakivubo; Water quality

INTRODUCTION

Heavy metal contamination of stream and river water ecosystem is a worldwide environmental problem. Trace amounts of heavy metals are always present in fresh waters from the weathering of rocks and soils (Muwanga, 1997; Anderson, 2003; Babel and Opiso, 2007; Samarghandi *et al.*, 2007; Igwe *et al.*, 2008; Al-Juboury, 2009). Recently, water quality monitoring has become a matter of concern in stream and river water systems affected by careless disposal of urban effluents. Runoff, atmospheric deposition and domestic and industrial effluent discharges are the major sources of aquatic pollution (Wasswa, 1997; Linnik and Zubenko, 2000; Campbell, 2001; Lwanga *et al.*, 2003 and Lomniczi *et al.*, 2007) and physico-chemical characteristics such as dissolved oxygen and the pH of aquatic ecosystems may determine

stream water ecosystem integrity. Monitoring of stream water physico-chemical characteristics and heavy metal concentration is necessary to establish the levels of contamination in wastewater. With increasing urbanisation and industrialization, there has been a rapid increase in industrial effluent discharge into the stream water, leading to increased pollution load. In aquatic eco-systems, trace elements may be immobilised within the stream water and may involve complex formation and co-precipitation as oxides and hydroxides of Fe, Mn or may occur in particulate form (Awofolu *et al.*, 2005; Mwiganga and Kansime, 2005; Nyangababo *et al.*, 2005b; Srivastava *et al.*, 2008).

The Nakivubo Channelized Stream is the major recipient of runoff and a surrogate end point for organic substances of industrial and domestic effluents (Kansime *et al.*, 1995; Sentongo, 1998). The Nakivubo channel is increasingly being polluted with direct

✉ *Corresponding Author Email: ssekaba@gmail.com
Tel.: +256 772 855 348; Fax: +256 414 501 974







with deionised water and again washed three times with the stream water before they were filled. For heavy metal analysis, a 500 mL bottle of linear polyethylene was used to collect water samples, filtered (using 0.45µm pore size) and three drops of concentrated nitric acid added for preservation. Also, 1 L bottle to collect water samples was used for determination of physico-chemical characteristics (total suspended solids and biological oxygen demand). All were stored in an ice box at 4°C and transported to Geology Laboratory at Makerere University and National Water and Sewerage Corporation central Laboratory at Bugolobi, respectively for analysis. Other physico-chemical characteristics such as dissolved oxygen, electrical conductivity, pH, temperature and total dissolved solids were determined on site using potable meters. The elements were then analyzed by direct aspiration of the sample solution into a Perkin-Elmer model 2380 flame atomic absorption spectrophotometer (AAS) after calibration with suitable elemental standards at close intervals.

Assessment of stream water contamination

Total suspended solids (TSS); Wastewater TSS classification consists of three classes (0-220 mg/L) ranging from weak to strong (< 100 mg/L) weak (100 – 220 mg/L) medium and strong wastewater (> 220 mg/L), [Akan et al \(2008\)](#).

Statistical analysis

Analysis of variance (ANOVA): ANOVA was employed

to determine whether groups of variables have the same means on data that are continuous or normally distributed and with homogeneous variance. Additionally, it was employed to assess the relationship between heavy metal concentrations and their elemental interaction between sections of the stream.

Correlation analysis: Pearson correlation's correlation analysis was adopted to analyse and establish inter-metal relationship and physico-chemical characteristics of the stream water.

Cluster analysis (CA) and Factor analysis (FA): CA was performed using Minitab Release 14 or 15 to classify elements of different sources on the basis of their similarities using dendrogram and to identify relatively homogeneous groups of variables with similar properties. FA was employed on the variables that are correlated to isolate or determine specific factors that are associated with such groupings of metal concentration to establish their origin. The data was standardized to give a normal distribution with a mean of 0 and a variance of 1. Sample means were standardized by subtracting the mean of their distribution and dividing by standard error (SE) or square root of the variance.

RESULTS AND DISCUSSION

Water chemical characteristics

Mean concentration of trace metals and physico-chemical parameters are given in [Table 2](#). The pH ranged from 6.30 to 9.83 for the Nakivubo Channelized



Fig. 2: Modified bulk sampler used in collection of rain water samples



Stream suggesting a slightly acidic to alkaline pH which fell within the range of surface water under natural conditions (5.0 to 8.6), the value recorded at 6th Street Bridge Mukwano fell well above the upper end of the range considered as normal, attributed to industrial wastewater outfall from Mukwano industries. Dissolved oxygen (DO) ranged from 0.00 to 6.69 mg/L Nakivubo Stream, 0.0 to 7.39 mg/L tributaries, 0.0 to 4.4 mg/L, industrial discharge outfall and 0.40 to 5.48 mg/L. Water was poorly aerated as indicated by very low dissolved oxygen in stream water which generally decreased downstream. WHO (2008) recommends 5.0 µm/L and above of DO but all stream water samples were below maximum permissible limit except Watindo Stream water which was relatively good. Dissolved oxygen below 5.0 mg/L may suggest aquatic contamination of such stream waters. Anoxic conditions are attributed to biochemical processes, oxidation of the substances, or decomposition of organic matter. Increased total dissolved solids (TDS) and biological oxygen demand (BOD) may suggest increased organic matter (OM) within the stream from industrial discharge, wastewater and effluents and ability of self-purification (Phiri *et al.*, 2005; Akan., *et al.*, 2007 and Akan *et al.*, 2008). City Abattoir (105-395 mg/L), peacock paint factory (14-2150 mg/L) and Sadolin paint factory/Meat parker outlet (214-3725 mg/L) are the biggest contributors of total suspended solids (TSS) and hence registered the highest BOD (5.6-1304 mg/L, 16.3-1806 mg/L and 137.2-3192 mg/L, respectively) and EC. This phenomenon indicated industrial emission into the Nakivubo Channelized Stream as the major source of contaminants.

The mean value of BOD at sites US01, US04, MS05, MS09, DS15, DS17, DST16 and all industrial discharge outlets to Nakivubo Channel exceeded the National Environment Management Authority (NEMA) critical level of 50.0 mg/L maximum permissible limits in effluent and wastewater (Table 2). Also, the mean value of total suspended solids at sites US01, US02, US04, MS05, DS15, CT03 and all industrial discharge outlets exceed the maximum permissible limit of 100 µm/L in wastewater or effluents and is classified as strong wastewater at Peacock paint factory, Sadoline paint factory and NWSC plant (> 220.0 mg/L) outfall. Electric conductivity (EC) was highest at industrial discharge outfall wastewater followed by tributaries and ranged between 397.0 and 1884.0 µs/cm and 208.0 to 961.0 µs/cm, respectively whereas sadoline paint factory ranged

between 208.0 and 961.0 µs/cm). EC showed a trend of gradual increase downstream, suggesting increased dissolved elements/substances to Murchison Bay into Lake Victoria. These results are consistent with those of Kar *et al.*, (2008). However, Total dissolved solids (TDS) were within National Electrical Manufacturers Association regulations for maximum permissible limit of discharge of wastewater and industrial effluents (1200 mg/L and 2000) (NEMA, 2006; WHO, 2008). Results of heavy metal concentration in stream water are shown in Table 3. Heavy metal concentrations in Nakivubo Channelized Stream water ranged from 44 ± 18 to 129 ± 70 µm/L Pb; 3 ± 1 to 9 ± 4 µm/L Cd; 8 ± 2 to 44 ± 35 µm/L Cu; 26 ± 6 to 59 ± 9 µm/L Zn; 505 ± 495 to 1035 ± 65 µm/L Mn and 300 ± 200 to 1450 ± 550 µm/L Fe. Stream water samples at Kiseka Market Bridge were the most contaminated by Pb, followed by the Luzira Culvert along the Nakivubo Channelized Stream. Generally, Pb concentration increased downstream but with the highest contributions from industrial discharge outfall.

The concentration of Pb exceeded the WHO (2008) for drinking water at most of the sites and NEMA, (2006) maximum permissible limit of discharge of wastewater into the environment (100 µm/L) at Kiseka Market, Kibira road tributary and Sadoline paint factory. However, Pb heavy metal concentrations at all sites exceeded 25 µm/L criterion for the protection of aquatic life. Total heavy metal content in stream waters along the tributaries to Nakivubo Channel ranged from 63 ± 24 to 106 ± 27 µm/L Pb; 3 ± 2 to 6 ± 2 µm/L Cd; 2 ± 4 to 8 ± 4 µm/L Cu and 51 ± 11 to 144 ± 24 µm/L Zn. Stream water along Kibira Stream tributary was the most contaminated with Zn, followed by Kitante stream and Lugogo. Lugogo stream water had the highest concentration of Cd (13 ± 6 µm/L) and Cu (18 ± 12 µm/L) followed by Kibira and Kayunga streams Cd (64 ± 20 µm/L) and Kitante Cu (16 ± 7 µm/L). The concentration of Cd exceeded WHO (2008) maximum permissible limit of 3 µm/L in wastewater. Heavy metal contents in stream waters at industrial discharge outlets range from 69 ± 25 to 113 ± 31 µm/L Pb; 4 ± 2 to 12 ± 5 µm/L Cd; 8 ± 3 to 18 ± 4 µm/L Cu; 51 ± 11 to 144 ± 25 µm/L Zn; 165 ± 35 to 2400 ± 700 µm/L and 600 ± 100 to 3050 ± 550 µm/L Fe. The range obtained from discharge outlets were within the NEMA standards for discharge of effluents and wastewater critical levels for Cd, Cu and Zn, except Pb which exceeds the limit of 100 µm/L as the maximum contaminant level at Sadoline paint



Table 2: Mean physicochemical characteristics of stream water samples from Nakivubo Channalized Stream, tributaries, industrial effluents and Watindo Stream (n=16)

Site	Temp (°C)			pHw			TDS (mg/L)			TSS (mg/L)			EC (µs/cm)			DO (mg/L)			BOD (mg/L)		
	Mean	Min.	Max.	Mean	Min.	Max.	Mean	Min	Max.	Mean	Min	Max.	Mean	Min.	Max.	Mean	Min.	Max.	Mean	Min	Max.
US1	26.93	22.90	30.20	7.11	6.30	7.83	235.06	215.00	271.00	176.13	16.00	850.00	470.88	432.00	544.00	2.51	0.48	4.28	62.33	11.20	176.90
US2	26.58	22.50	30.10	7.07	6.55	7.97	178.81	145.00	255.00	118.31	5.00	733.00	356.75	299.00	502.00	3.32	0.00	6.69	36.08	5.60	188.60
US3	27.79	23.40	31.20	7.28	6.92	7.99	236.88	187.00	359.00	82.81	19.00	185.00	477.13	377.00	763.00	2.55	0.63	4.72	44.54	10.92	145.60
US4	27.51	23.30	30.30	7.18	6.30	7.89	251.00	114.00	298.00	195.13	48.00	357.00	471.52	5.30	618.00	1.80	0.00	3.26	166.97	58.52	952.00
MS5	27.13	24.10	29.60	7.19	6.84	7.94	298.66	3.57	433.00	278.50	109.00	1400.00	596.32	7.16	867.00	1.40	0.00	3.96	136.90	11.20	351.68
MS9	25.79	21.80	28.30	7.78	6.94	9.83	264.56	178.00	463.00	54.44	12.00	161.00	534.75	344.00	923.00	1.48	0.00	3.50	83.59	7.70	672.00
DS15	24.49	22.50	26.10	7.55	7.17	8.22	357.94	190.00	443.00	172.94	29.00	410.00	736.99	540.00	892.00	0.77	0.00	2.08	140.43	7.14	560.00
DS17	22.87	21.40	24.80	7.05	6.70	7.89	334.69	236.00	426.00	86.59	9.40	194.00	667.31	473.00	807.00	1.02	0.00	2.15	71.75	23.52	280.00
MT7	24.98	21.80	27.40	7.33	6.91	7.86	305.50	236.00	361.00	69.19	9.00	367.00	612.63	479.00	722.00	1.62	0.00	3.98	49.09	6.24	200.00
MT10	25.36	23.30	27.40	7.09	6.59	9.35	129.69	102.00	166.00	24.19	6.00	49.00	260.63	208.00	335.00	3.68	1.20	7.39	30.31	5.18	156.20
MT13	22.97	21.50	25.10	7.15	6.35	7.62	199.81	163.00	478.00	24.13	3.00	128.00	405.38	328.00	961.00	3.23	0.30	6.42	46.66	5.60	224.00
DT16	24.71	22.90	27.20	6.86	6.12	9.36	263.56	156.00	405.00	115.31	18.00	289.00	526.06	319.00	807.00	1.86	0.00	4.31	155.87	12.46	672.00
MD8	27.27	22.90	40.20	7.53	7.01	9.64	352.47	281.00	529.00	141.67	16.00	740.00	704.00	570.00	1024.00	1.71	0.00	4.40	124.94	16.10	856.80
MD11	24.76	21.50	27.80	6.97	6.45	7.62	513.38	224.00	935.00	404.50	14.00	2150.00	1037.81	449.00	1884.00	2.71	1.22	3.99	255.12	16.25	1806.00
MD12	24.71	23.00	26.00	7.43	6.99	8.08	512.19	220.00	1200.00	1243.19	214.00	3725.00	955.31	488.00	1316.00	0.98	0.00	2.45	1337.51	137.20	3192.00
MD14	25.27	23.00	27.50	7.61	7.15	8.31	600.50	199.00	747.00	237.44	105.00	395.00	1198.56	397.00	1500.00	1.47	0.00	3.82	251.60	5.60	1304.80
CT1	25.04	20.00	31.60	7.07	6.61	7.84	83.90	35.00	575.00	79.31	15.00	555.00	104.34	80.40	125.10	4.09	0.60	5.48	41.91	8.40	298.10
CT2	24.93	20.30	32.00	7.00	6.55	7.72	51.83	40.00	60.80	97.97	12.00	877.50	103.11	80.10	122.20	4.21	3.13	5.39	16.69	2.24	38.08
CT3	25.06	19.30	35.90	6.88	6.51	7.51	53.19	12.80	69.60	105.87	12.00	370.00	106.23	25.70	140.20	3.66	0.40	5.24	31.97	5.60	197.10
**WHO	40						2000						1000			5.0			50.0		
**NEMA	20-35			6-8			1200			100						5.0			50.0		

**WHO, (2008) maximum permissible limit of wastewater into aquatic habitat

***National Environmental Management Authority, (2006/2007) maximum permissible limit of wastewater into aquatic habitat; CT1, CT2 and CT3 are sites along Watindo Stream



factory discharge outlet, Kisekka market and Kibira Road Stream which is a surrogate discharge outlet for Uganda batteries Limited factory and Nice House of plastics factory (Table 3). Heavy metal concentration in rain for Pb (13 µm/L and 33 µm/L) was above water quality criterion value set for domestic supplies at WHO standards whereas, the rest were below the maximum permissible limits. The levels of concentrations of lead in rain water were far below that in stream water. Elemental concentrations of Pb, Cd, Cu and Zn were below the concentrations in effluents observed by Nabulo *et al.*, (2008) except Cd at Peacock paint factory effluent which was in the same order of magnitude. However heavy metal concentrations in stream water along the Nakivubo Channel were in the same order of magnitude with those observed by Muwanga and Balifaijo, (2006).

Manganese concentrations exceeded the limit of 1000 µm/L as the maximum permissible limit of wastewater discharge into the environment at

Mukwano industries, Peacock paint factory outfall and Kayunga tributary inflows leading to its accumulation downstream. The concentration of heavy metals along Watindo Stream ranged from 50 ± 18 to 106 ± 47 µm/L Pb; 2 ± 1 to 5 ± 2 µm/L Cd; 5 ± 2 to 14 ± 6 µm/L Cu; 20 ± 4 to 241 ± 5 µm/L Zn; 155 ± 145 to 165 ± 135µm/L Mn and 4050 ± 250 to 4600 ± 1000 µm/L Fe (Table 3). The mean Cd concentration in stream water at site CT01 and CT02 exceeded WHO (2008) maximum permissible limit for drinking water. But Pb, Cu, Mn and Fe concentrations in water of several magnitudes higher than surface water standards (Table 3). This phenomenon is of common occurrence in most populated urban areas close to industrial establishments (Mohiuddin *et al.*, 2010).

Analysis of variance: The two-way ANOVA in table 4a-b indicate that sites show no significant effect on variation between groups of means of the heavy metals at deferent stream sections (Table 4a). However, the mean heavy metal concentrations of elements within

Table 3: Mean values and SE of trace metal concentrations in water at different sites along Nakivubo Channelized Stream and its Tributaries, Industrial effluent, rain water and Watindo Stream (n=16)

Sites	Heavy metals (µg/L)					
	Pb	Cd	Cu	Zn	Mn	Fe
Nakivubo stream						
US01	44±18	4±2	9±30	26±6	600±500	300±300
US02	45±20	4±2	12±6.0	26±5	535±465	500±100
US03	129±70	9±4	44±35	34±7	505±495	300±200
US04	3±9.0	3±1	8±2.0	46±7	685±215	900±400
MS05	53±26	5±2	27±20	59±9	955±145	1250±350
MS09	63±24	4±2	21±12	37±6	1000±300	1200±100
DS15	81±23	6±4	15±6	51±8	675±125	1050±350
DS17	88±24	4±2	13±6	33±5	1035±650	1450±550
Tributaries						
MT07	64±20	4±2	6±2	31±5	1915±185	2000±700
MTS10	63±24	3±2	16±7	49±14	155±145	250±250
MT13	75±30	3±6	18±12	32±6	475±425	1000±200
DT16	106±27	4±2	9±3	157±34	475±25	1250±450
Industrial Effluents						
MD08	100±34	4±2	25±0.008	65±18	1415±1384	900±800
MDS11	69±25	12±5	8±0.003	51±11	2400±700	800±700
MD12	113±31	6±2	18±0.004	144±24	670±30	3050±550
MD14	93±37	3±2	15±0.003	92±15	165±35	600±100
Watindo stream						
CT01	106±47	4±2	14±0.006	20±4	160±140	4050±250
CT02	75±31	5±2	5±5	24±5	155±145	4300±400
CT03	50±18	2±1	11±6	22±4	165±135	4600±100
Rain water						
Rw1n	13±9	2±1	14±7	18±5	50±50	200±200
Rw2w	33±19	3±1	25±20	21±3	10±10	100±0
**WHO, 2008	10	3	2000	3000	100	300
* NEMA, 2006	100	100	1000	5000	1000	
***surface water	0.015	-	1.3	-	0.05	0.3

* National Environment Management Authority maximum permissible limit of wastewater into aquatic habitat; ** World Health Organisation maximum permissible limit of drinking water; ***Wedepohl, (1989) for surface water; Rw1n: rainwater collected in experimental site (Nakivubo); Rw2w: rainwater collected in control site (Namalere research centre-Watindo)



sites differed significantly at different stream sections; upstream ($F_{3,15}=27.33$, $P < 0.001$); midstream ($F_{1,5}=1828.280$, $P < 0.001$) and downstream ($F_{1,5}=51.620$, $P < 0.001$), at $P = 0.05$. Sections of the stream have significant effect on the mean heavy metal concentrations ($F_{2,30}=13.3$, $P < 0.001$), Table 4b. However, there was also a significant difference among heavy metal concentrations between sections of the stream ($F_{2,30}=139.26$, $P < 0.001$). The effects of interaction between sections and mean heavy metal concentration was significant ($F_{2,30}=5.79$, $P < 0.001$) suggesting a transport phenomenon or mobility of heavy metals.

Correlation analysis: Pearson correlation coefficients for the data were evaluated to determine the level of inter-metal association and the values are shown in Table 5. Elemental pairs Pb/Cd ($r = 0.56$); Pb/Zn ($r = 0.50$); Cd/Mn ($r = 0.52$) and Mn/Fe ($r = 0.56$) were significantly correlated with each other, suggesting that each paired elements have identical source or chemical phenomena, whereas the rest of the elemental pairs are not correlated with each other. The association of Mn with Fe in stream water may strengthen terrigenous sources and/or geo-chemistry. Metals show correlation with the physico-chemical characteristics. Pairs Pb/TDS, Pb/EC, Cd/EC, Zn/EC, Zn/BOD and Mn/EC are positively correlated with each other and Zn/DO and Fe/DO were negatively correlated, suggesting their Physico-chemical associations, whereas the rest of the metals and physico-chemical parameters are not

correlated. Positive correlation of Cd/EC elemental pair was consistent with that of Kar *et al.*, (2008). Table 5 also shows that pairs BOD/TDS, BOD/TSS, DO/EC, TDS/DO, EC/TDS and TSS/TDS were significantly correlated with each other, whereas the rest of the pairs are not correlated. Very high BOD and very low DO in the stream water suggest pollution by organic matter and decomposition as a self purification mechanism. This phenomenon results into a reducing condition in organic rich stream water. These results are in line with those done by Igbiosa and Okoh, (2009). The development of reducing condition in stream water promotes the reduction of iron and manganese hydroxides (Linnik and Zubenko, 2000). Under slightly acidic to alkaline condition, Cd specifically adsorbed proportion is high in solid phase and Pb may occur bonded to poorly crystalline oxides (Fergusson 1990). The pH values (slightly acidic to alkaline), may limit the degree of desorption of heavy metals into solution and can promote co-precipitation except Pb at $pH > 6$ the solubility is governed by soluble organic ligands under reducing conditions (Muwanga, 1997). Contaminants released to flow in restricted water bodies at the mixing event can promote co-precipitation (Anazawa *et al.*, 2004).

Cluster analysis and factor analysis: Cluster analysis was performed on the data using Average linkage and correlation coefficient distance. Results of cluster analysis are shown in Fig. 3. Four groups of elemental associations with eigenvalue > 1 were

Table 4a: Two-way ANOVA results for sites and heavy metal concentration variables (Dependant variables were log-normal transformed)

Stream sections	Source of Variation	DF	SS	MS	F	P
Upstream	Sites	3	0.003	0.001	0.920	0.454
	Metal content	5	0.162	0.032	27.330	0.000
Midstream	Sites	1	0.000	0.000	0.180	0.689
	Metal content	5	0.255	0.051	1828.280	0.000
Downstream	Sites	1	0.002	0.002	2.210	0.197
	Metal content	5	0.238	0.048	51.620	0.000

DF-degree of freedom; F-factor mean square SS- Sum of squares; MS- Mean square; P = 0.05

Table 4b: Two-way ANOVA results for stream sections and mean heavy metal concentration variables (dependant variables were log-normal transformed)

Source of Variation	DF	SS	MS	F	P
Sections	2	0.000	0.01232	13.3	0.000
Metal content	5	1.000	0.12904	139.26	0.000
Sections* Metal	10	0.000	0.00536	5.79	0.000

SS- Sum of squares; MS- Mean square; P = 0.05



extracted in the analysis. Heavy metal elements were fused into clusters because of their relative elemental concentrations at each site and their similarity coefficients. Group I contains Pb and Cu while group II consists of Cd, Mn and Fe. Group III contains Zn as well as BOD, TDS, EC, TSS and stream water pH.

Biplot of sites and elemental concentrations suggest that Kibira Road stream is a source of Zn, Pb and copper, whereas Mukwano industries is a point source of Zn and Pb (Fig. 3). Sadoline paint factory was the point source of Pb and Zn, whereas NWSC is the point source of Cu, Zn and Pb. Inflows of Kayunga Stream into the Nakivubo Channelized Stream comes with Mn and Cd. The lengths of the arrow for Pb and Zn suggest that NWSC were the highest source, Peacock paint factory for Fe, Kayunga and Lugogo streams for Mn and Cd.

The first factor accounts for 35.7 % of the total variance and contains Zn and Fe as well as water pH, total dissolved solids, TSS, electrical conductivity and BOD, with high variable loadings on this factor (Table 6) and corresponds to group III of the cluster analysis. Organic factor may indicate point sources such as municipal and industrial

effluents and adsorption by TDS as well as flocculation or co-precipitation which is pH controlled (Balachandran *et al.*, 2005; Karbassi *et al.*, 2007; Abdel-Ghani and Elchaghaby, 2007; Abdel-Ghani *et al.*, 2009) TDS and TSS seems to play a major role in the adsorption of heavy metals in stream water rich with organic matter. These phenomena remove heavy metals rapidly and render it free of contamination under varying dissolved salts.

The second factor accounts for 19.9 % of the variance and contains Cd, Mn and Fe with high variable loadings on this factor which corresponds to group II of the cluster analysis. This indicates co-precipitation of Cd with Mn and Fe oxides/hydroxides. The third factor accounts for 16.8 % of the variance and contains Pb, Cd and Cu with high variable loadings on this factor and corresponds to group I of the cluster analysis. This factor is considered to represent anthropogenic toxic pollutants from metal works and fabrications, vehicular and industrial emissions. Distribution of Zn between different factors 1 and 2 may suggest dual sources whereas pH indicates its influence as a controlling factor on heavy metal co-precipitation.

Table 5: Pearson correlation coefficient matrix (r) for mean heavy metal contents and physicochemical characteristics in Nakivubo Channelized Stream water (n=16)

	Pb	Cd	Cu	Zn	Mn	Fe	Temp	pHw	TDS	TSS	EC	DO	BOD
Pb	1												
Cd	0.56*	1											
Cu	0.30	0.12	1										
Zn	0.50*	0.14	-0.01	1									
Mn	0.23	0.52*	-0.09	-0.18	1								
Fe	0.27	0.33	-0.19	0.38	0.56*	1							
Temp	-0.35	-0.29	-0.07	-0.01	-0.11	-0.34	1						
pHw	0.25	-0.18	0.25	0.36	-0.20	0.00	0.11	1					
TDS	0.49*	0.32	-0.17	0.47	0.31	0.48*	-0.20	0.42	1				
TSS	-0.13	-0.02	-0.28	0.13	-0.05	0.14	0.21	0.17	0.56*	1			
EC	0.67**	0.48*	-0.14	0.55*	0.49*	0.43	-0.14	0.30	0.83**	0.18	1		
DO	-0.54	-0.22	0.03	-0.53*	-0.44	-0.56*	0.26	-0.42	-0.77**	-0.25	-0.83**	1	
BOD	-0.07	-0.14	-0.14	0.53*	-0.24	0.44	-0.10	0.31	0.48*	0.65**	0.13	-0.32	1
pHs	0.12	0.36	0.12	0.03	0.35	0.15	0.49*	-0.03	-0.21	-0.004	0.03	0.02	-0.20

* Correlation is significant at the 0.05 level (2-tailed), ** Correlation is significant at the 0.01 level (2-tailed) pHw; water pH, pHs; sediment pH






Fig. 3: Dendrogram of elemental concentrations and physico-chemical parameters of Nakivubo Channelized stream water and industrial discharge effluents (n=16)




Fig. 4: Biplot of elemental concentrations (a) and sites (b) of stream water



Table 6: Rotated factor loadings and communalities of elements and physico-chemical characteristics in stream water samples (n=16)

Variable	Factor 1	Factor 2	Factor 3	Communality
Pb	0.152	0.075	0.865	0.777
Cd	0.121	-0.613	0.525	0.666
Cu	-0.237	0.122	0.791	0.697
Zn	0.451	0.608	0.211	0.618
Mn	0.102	-0.896	-0.044	0.815
Fe	0.614	-0.428	0.077	0.567
pH-w	0.352	0.589	0.266	0.542
TDS	0.924	0.041	0.068	0.859
TSS	0.799	0.036	-0.231	0.693
EC	0.944	0.041	0.27	0.909
BOD	0.858	0.294	-0.045	0.824
Variance	3.9269	2.1912	1.8494	7.9675
% Var.	0.357	0.199	0.168	0.724

CONCLUSION

The Nakivubo Stream water is stripped of heavy metal contaminants. Because of this, the quality of the Nakivubo aquatic ecosystem is improved. However, high anoxic conditions attributed to biochemical processes and decomposition of organic matter as a purification mechanism (organic matter pollution) poses a threat to aquatic life. The greatest threat in Nakivubo Channelized Stream water would be posed by Pb and Cd which almost always occur as mobile elements in water under neutral to alkaline conditions but are always rapidly removed. Flocculation and co-precipitation of Mn and Fe hydroxides, water pH and TDS could be the major mechanism explaining the rapid removal of Pb and Cd to bottom sediments. The levels of dissolved oxygen were below the maximum permissible limit, while the mean biological oxygen demand was above the maximum permissible limit. FA results reveal two sources of pollutants as explained by three factors (72.40 %); (i) mixed origin or chemical phenomena of industrial and vehicular emissions, and (ii) dual origin of Zn (vehicular, commercial establishment and industrial). Atmospheric deposition of heavy metals especially Pb in rain is apparent and may be attributed to vehicular emissions.

ACKNOWLEDGEMENTS

The authors are thankful to Kampala International University for the financial support in form of PhD research project and Department of Geology, Faculty

of science, Makerere University, for geochemical analyses.

REFERENCES

- Abdel-Ghani, N. T.; Elchaghaby, G. A., (2007). Influence of operating conditions on the removal of Cu, Zn, Cd and Pb ions from wastewater by adsorption. *Int. J. Environ. Sci. Tech.*, 4 (4), 451-456 (6 pages).
- Abdel-Ghani, N. T.; Hegazy, A. K.; El-Chaghaby, G. A., (2009). Typha domingensis leaf powder for decontamination of aluminium, iron, zinc and lead: Biosorption kinetics and equilibrium modeling. *Int. J. Environ. Sci. Tech.*, 6 (2), 243-248 (6 pages).
- Akan, J. C.; Abdulrahman, F. I.; Dimari, G. A.; Ogugbuaja, V. O., (2008). Physicochemical determination of pollutants in wastewater and Vegetable samples along the Jakara Wastewater Channel in Kano Metropolis Kano State, Nigeria. *Europ. J. Sci. Res.*, 23 (1), 122-133 (12 pages).
- Akan, J. C.; Abdulrahman, F. I.; Ayodele, J. T.; Ogugbuaja, V.O., (2007). Studies on the effect of Municipal waste and Industrial effluents on the Pollutant level of River Challawa, Kano State, Nigeria. *Res. J. Appl. Sci.*, 2 (4), 530-535 (6 pages).
- Al-Juboury, A. I., (2009). Natural Pollution By Some Heavy Metals in the Tigris River, Northern Iraq. *Int. J. Environ. Res.*, 3 (2), 189-198 (10 pages).
- Anderson, D., (2003). Introduction to heavy metal monitoring. UK.
- Anazawa, K.; Kaida, Y.; Tomiyasu, T.; Sakamoto, H., (2004). Heavy metal distribution in river waters and sediments around a "Firely Village", Shikoku, Japan: Application of Multivariate analysis. *Japan Soc. Anal. Chem.*, 20 (1), 79-84 (6 pages).
- Awofolu, O. R.; Mbolekwa, Z.; Mtshemla, V.; Fatoki, O. S., (2005). Levels of heavy metals in water and sediments from Tyume River and its effects on an irrigated farmland. *Water SA*. 31 (1), 87-94 (8 pages).
- Babel, S.; Opiso, E. M., (2007). Removal of Cr from synthetic wastewater by sorption into volcanic ash soil. *Int. J. Environ. Sci. Tech.*, 4 (1), 99-107 (9 pages).
- Balachandran, K. K.; Lalu Raj, C. M.; Nair, M.; Joseph, T.; Sheeba, P.; Venugopal, P., (2005). Heavy metal accumulation in a flow restricted, tropical estuary. *Estuar. Coast. Shelf Sci.*, 65 (1-2), 361-370 (10 pages).
- Campbell, L. M., (2001). Mercury in Lake Victoria (East Africa): Another emerging issue for a Beleaguered Lake? PhD, Thesis. Waterloo, Ontario, Canada.
- Fergusson, J. E., (1990). The heavy elements: Chemistry, environmental impact and health effects. Pergamon Press, Oxford, 614.
- Igbinosa, E. O.; Okoh, A. I., (2009). Impact of discharge wastewater effluents on the physico-chemical qualities of a receiving water shade in typical rural community. *Int. J. Environ. Sci. Tech.* 6 (2), 175-182 (8 pages).
- Igwe, J. Ch.; Abia, A. A.; Ibe, Ch. A., (2008). Adsorption kinetics and intraparticle diffusivities. *Int. J. Environ. Sci. Tech.*, 5 (1), 83-92 (10 pages).
- Kansiime, F.; Katayo, E.; Okot-okumu, J., (1995). Effects of pollution on Inner Murchison bay (Lake Victoria-Uganda) on the distribution and abundance of plankton. A



- report of Makerere Institute of Environment and Natural Resources.
- Kar, D.; Sur, P.; Mandal, S. K.; Saha, T.; Kole, R. K., (2008). Assessment of heavy metal pollution in surface water. *Int. J. Environ. Sci. Tech.*, 5 (1), 119-124 (6 pages).
- Karbassi, A. R.; Nouri, J.; Ayaz, G. O., (2007). Flocculation of trace metals during mixing of Talar river water with Caspian Seawater. *Int. J. Environ. Res.*, 1 (1), 66-73 (8 pages).
- Linnik, P. M.; Zubenko, I. B., (2000). Role of bottom sediments in the secondary pollution of aquatic environments by heavy metal compounds. *Lakes and Reservoirs. Res. Manag.*, 5 (1), 11-21 (10 pages).
- Lomniczi, I.; Boemo, A.; Musso, H., (2007). Location and characterisation of pollution sites by principal component analysis of trace contaminants in a slightly polluted seasonal river: A case study of the Arenales River (Salta, Argentina). *Water SA.*, 33 (4), 479-485 (7 pages).
- Lwanga, M. S.; Kansime, F.; Denny, P.; Scullion, J., (2003). Heavy metals in Lake George, Uganda with relation to metal concentrations in tissues of common fish species. *Hydrobiology*, 499 (1-3), 83-93 (11 pages).
- Muwanga, A., (1997). Environmental impacts of copper mining at Kilembe, Uganda: A geochemical investigation of heavy metal pollution of drainage waters, stream, sediments and soils in the Kilembe valley in relation to mine waste disposal. Ph.D. Dissertation, Universitat Braunschweig, Germany.
- Muwanga, M.; Balifajjo, E., (2006). Impact of industrial activities on heavy metal loading and their physiochemical effects on wetlands of Lake Victoria Basin (Uganda). *African J. Sci. Tech.*, 7 (1), 51-63 (13 pages).
- Mohiuddin, K. M.; Zakir, H. M.; Otomo, K.; Sharmin, S.; Shikazono, N., (2010). Geochemical distribution of trace metal pollutants in water and sediments of downstream of an urban river. *Int. J. Environ. Sci. Tech.*, 7 (1), 17-28 (12 pages).
- Mwiganga, M.; Kansime F., (2005). Impact of Mpererwe landfill in Kampala Uganda, on the surrounding environment. *Phys. Chem. Earth.*, 38 (11-16), 91-95 (5 pages).
- Nabulo, G.; Oryem Origa, H.; Nasinyama, G. W.; Cole, D., (2008). Assessment of Zn, Cu, Pb and Ni contamination in wetland soils and plants in the Lake Victoria basin. *Int. J. Environ. Sci. Tech.*, 5 (1), 65-74 (10 pages).
- NEMA (2006). National Environment Management Authority., State of environment report for Uganda.
- Nyangababo, J. T.; Henry, I.; Omutunge, E., (2005a). Heavy metal contamination in plants, sediments and air precipitation of Katonga, Simiyu and Nyando wetlands of Lake Victoria basin, East Africa. *Bull Environ. Contam. Toxicol.*, 75 (1), 189-196 (8 pages).
- Nyangababo, J. T.; Henry, E.; Omutunge, E., (2005b). Lead, cadmium, copper, manganese and zinc in wetland waters of Victoria lake basin, East Africa. *Bull. Environ. Contam. Toxicol.*, 74 (5), 1003-1010 (8 pages).
- Phiri, O.; Mumba, P.; Moyo, B. H.; Kadewa, W., (2005). Assessment of the impact of industrial effluent on water quality of receiving rivers in urban areas of Malawi. *Int. J. Environ. Sci. Tech.*, 2 (3), 237-244 (8 pages).
- Samarghandi, M. R.; Nouri, J.; Mesdaghinia, A. R.; Mahvi, A. H.; Nasser, S.; Vaezi, F., (2007). Efficiency removal of phenol, lead and cadmium by means of UV/TiO₂/H₂O₂ processes. *Int. J. Environ. Sci. Tech.*, 4 (1), 19-25 (7 pages).
- Sentongo, J., (1998). Assessment of pollution to Lake Victoria by industrial and municipal activities around Lake Victoria in Uganda., M.Sc. Thesis., Makerere University, Uganda.
- Srivastava, A.; Gupta, S.; Jain, V. K., (2008). Source apportionment of suspended particulate matter in coarse and fine size ranges over Delhi. *Aerosol Air Qual. Res.*, 8 (2), 188 – 200 (13 pages).
- Wasswa, J., (1997). Determination of levels of pollutants within sediments in water channels from industries in selected parts of Lake Victoria. M.Sc., Thesis, Makerere University, Uganda.
- WHO., (2008). Guidelines for drinking - water Quality 3rd Edition incorporating the first and second agenda volume 1 Recommendations, World Health Organizations Geneva.

AUTHOR (S) BIOSKETCHES

Sekabira, K., Ph.D. Scholar, Department of Environment, Faculty of Science and Engineering, Kampala International University, P. O. Box 20000, Kampala, Uganda; Email: ssekaba@gmail.com

Oryem Origa, H., Ph.D., Professor, Department of Botany, Faculty of Science, Makerere University, P. O. Box 7062, Kampala, Uganda. Email: horyem_origa@botany.mak.ac.ug

Basamba, T. A., Ph.D., Department of Soil Science, Faculty of Agriculture, Makerere University, P. O. Box 7062, Kampala, Uganda. Email: ateenyitwaha@hotmail.com

Mutumba, G., Ph.D., Assistant Professor, Department of Botany, Faculty of Science, Makerere University, P. O. Box 7062, Kampala, Uganda. Email: gmutumba@botany.mak.ac.ug

Kakudidi, E., Ph.D., Department of Botany, Faculty of Science, Makerere University, P. O. Box 7062, Kampala, Uganda. Email: ekakudidi@sci.mak.ac.ug

How to cite this article: (Harvard style)

Sekabira, K.; Oryem Origa, H.; Basamba, T. A.; Mutumba, G.; Kakudidi, E., (2010). Heavy metal assessment and water quality values in urban stream and rain water. *Int. J. Environ. Sci. Tech.*, 7 (4), 759-770.

